



FIFTY SHADES OF PURPLE: Immediate Treatment of Ankle Sprains

The very common ankle sprain can be more complex than you think and requires proper management to ensure a full recovery. The following leaflet outlines what you should do immediately after you sprain your ankle. Further leaflets in the series outline exercises you can do to help in the longer term rehabilitation of the injury.

WHAT IS AN ANKLE SPRAIN?

An ankle sprain is a stretch or tear in one or more of the lateral (outside) ligaments of the ankle. Ankle ligaments are slightly elastic bands of tissue that keep the ankle bones in place. Because the ankle is responsible for both weight-bearing and mobility, it is particularly susceptible to injury. The relatively small joint has to withstand large forces exerted when walking, running and jumping, especially if the surface is uneven.

Most ankle sprains happen when the ankle twists or rolls suddenly, usually a rapid and uncontrolled movement. The most common injuries happen when the foot rolls onto the outside of the ankle, straining the outside ligaments of the ankle joint. Symptoms of a sprained ankle include; pain, tenderness and swelling, bruising, trouble moving the ankle, and sometimes an inability to put your full weight on the ankle.

HOW LONG DOES IT TAKE TO HEAL?

Most people recover completely from mild sprains within two to six weeks. More severe sprains can take up to six months before you can return to full activity, or sport. Once a significant sprain occurs, without good rehabilitation the joint may never be as strong as it was before the injury. It is not surprising therefore that many people have a history of repeated ankle sprains. With the correct rehabilitation however, you can help your ankle become even stronger than it was before the injury.

WHAT CAN BE DONE?

Tissue injury usually involves damage to small blood vessels that results in bleeding at the site of injury. This bleeding leads to inflammation, part of the natural healing process. However, the body tends to overreact to sudden traumatic injury and as a result excess inflammatory fluid accumulates which can result in 'scar' tissue production. Too much scar tissue may prevent normal function with reduced flexibility and increased risk of re-injury.

It is important to get medical advice to gain a positive diagnosis and

correct treatment. Follow the PRICE guidelines immediately after injury and for at least 3 days afterwards before doing anything else. Your local physical therapist as well as most massage therapists can assist you with this:

PROTECT - Protect the injured tissue from undue stress and avoid ALL movements in the same direction as when the injury occurred.

REST - Unload the joint (take the weight off it) as much as you can in the first 72 hours after injury. Try and avoid walking on the joint as much as possible.

ICE - Ice is an amazing natural healer and a great short-term pain reducer. It is also believed to have a beneficial effect in reducing swelling and promoting healing. The optimal amount of time to apply ice is around 10-15 minutes in bony areas such as the ankle. It can be applied as often as desired to achieve pain relief, ideally every 1-2 hours.

COMPRESSION - This is advised for the first 72 hours, but only while your foot *isn't* elevated. The compression can be firm as long as it doesn't cause pins and needles or any loss of feeling around the joint.

ELEVATION - Reduces the flow of blood to the area which helps reduce swelling. Elevation is definitely recommended in the first 72 hours after injury. However remove any compression while your foot is elevated unless you are wearing just a light compression bandage.

When following **PRICE** it is also important to avoid **HARM**, hence the saying: '*Give PRICE and avoid HARM*'. HARM is an acronym for Heat, Alcohol, Running, Massage.

Following this acute management phase, your physical therapist will start some 'hands-on' treatment to mobilise and strengthen the joint. This phase of treatment is crucial to ensure you return to full function and prevent future injury.

Adequate preparation for activity is key and weight-bearing should progress gently. Drastic changes in activity level and performing unpractised skills expose your ankle to re-injury. Gradually build up your fitness level. Your ankle, and the rest of your body, will thank you for it!

The information contained in this article is intended as general guidance and information only and should not be relied upon as a basis for planning individual medical care or as a substitute for specialist medical advice in each individual case. ©Co-Kinetic 2017